

RSV Products Visual Guide

FOR PATIENTS WHO ARE: Pregnant

Administer **ABRYSVO (Respiratory Syncytial Virus Vaccine)**. Abrysvo is supplied in a kit that includes a vial of Lyophilized Antigen Component, a prefilled syringe containing diluent, and a vial adapter.

Timing: A single dose during weeks 32 through 36 of pregnancy from September through January in most of the U.S.



FOR PATIENTS WHO ARE:

A newborn or infant*

Administer **BEYFORTUS (nirsevimab-alip)**. Beyfortus is supplied in a prefilled syringe. 50mg doses are light blue and 100mg doses are purple.

Timing: A single dose just before or during the RSV season (from October through the end of March in most of the U.S.).



Premature/at high risk*

Administer **SYNAGIS (Palivizumab)**. Synagis is supplied in single-use vials. 50mg vials are pink and 100mg vials are blue.

Timing: One injection given monthly for up to 5 months throughout the RSV season.



FOR PATIENTS WHO ARE: 60 years or older**

Administer **AREXVY (Respiratory Syncytial Virus Vaccine, Adjuvanted)**, **ABRYSVO**, or **mRESVIA (Respiratory Syncytial Virus Vaccine)**.

Timing: A single dose any time, ideal timing is from August through October. *A one time vaccination.*



*See full clinical guidance for additional prescribing information, age cut-offs, special circumstances, etc.

**People ages 60–74 should only be vaccinated against RSV if they are at increased risk of severe RSV, meaning they have certain chronic medical conditions, such as lung or heart disease, or they live in nursing homes or other long-term care facilities. Everyone ages 75 and older should be vaccinated.